

The Effect of Ankle Range of Motion on Balance Performance of Elderly People

JEONG-WOO LEE, PhD¹⁾, SE-WON YOON, PhD¹⁾, JUNG-HWAN KIM, MS¹⁾,
YOUNG-PIL KIM, MS²⁾, YONG-NAM KIM, PhD³⁾

¹⁾ Department of Physical Therapy, Kwangju Women's University

²⁾ Best & Hydrotech Institute

³⁾ Department of Physical Therapy, Nambu University: 864-1 Wolgye dong, Gwangsan-gu, Gwangju, 506-706 Republic of Korea. TEL: +82 2-62-970-0231, FAX: +82 2-62-0118, E-mail: kyn0231@nambu.ac.kr

Abstract. [Purpose] This study examined the influence of range of motion of the ankle joints on elderly people's balance ability. [Methods] We conducted a four-week experiment using 60 out of the 89 elderly people who used B senior health facility, as subjects. TETRAX stability, synchronization were conducted to measure balance ability. We examined correlations between TETRAX balance and ankle joint ROM. Student's t-test and Pearson correlation coefficients were used for the statistical analyses. [Results] There were significant correlations between measurement items of TETRAX stability and synchronization and movement of the ankle joint. [Conclusion] It was found that as had plate was stable in measurement using ankle joint ROM and TETRAX after closing eyes, balance was maintained with Dorsi Flexion.

Key words: Ankle range of motion, Balance, TETRAX

(This article was submitted May 7, 2012, and was accepted May 29, 2012)

INTRODUCTION

The elderly often fall because their muscle mass, muscle strength and range of motion reduce as we grow older. Walking speed and pace also decrease as well as balance ability reduced¹⁾. Falls by the elderly are significantly associated with decline in their ability to adjust posture²⁾, dynamic balance ability³⁾, walking speed and mobility⁴⁾, and muscle strength in the lower extremity⁵⁾. Also, the elderly experience falls due to decrease of muscle strength in the lower extremity, balance and flexibility^{6, 7)}. Thus, a decline in balance and functional mobility increases the risk of falling⁸⁾, and balance ability is one of the important factors related to falls.

Posture and balance adjustment is achieved by integration of information from the visual, vestibular and somatosensory systems⁹⁾. Lord et al.¹⁰⁾ measured the balance ability of elderly people, and reported that there are high correlations among hypofunction of the somatosensory system, especially proprioception and balance ability, and that the visual and vestibular systems are subsidiary factors for maintenance of balance. Center of pressure (COP), which is an important factor for maintaining balance, is defined as the vertical component of force at the body's point of contact with the ground. It is the same as the vertical force of the center of gravity (COG)¹¹⁾, and if the center of pressure is outside the base of support, falls or staggering is caused¹²⁾.

A Hindrance factor such as unstable ground in maintaining balance can be postural sway¹³⁾. According to Teasdale et al.¹⁴⁾, the decline of stability as measured by postural sway is

caused by decrease of sensitivity in the visual and vestibular systems and proprioception, and postural sway increases as we grow older¹⁵⁾. Also, as we grow older, decrease of visual function and range of motion in the lower extremities may be factors impairing balance ability¹⁶⁾.

The ankle joints have important roles in maintaining balance¹⁷⁾. All movements of ankle joints are related to the maintainance of in walking, and they modulate interaction between foot and base. They are vital for walking and balance¹⁸⁾, and are used to support the limbs and adjust posture¹⁹⁾. Ankles, muscles, and joints of the feet are designed to provide stability as well as mobility in the terminal structure of lower extremities²⁰⁾.

Since the range of motion tends to be decrease with the dynamic characteristics of joint morphology as we grow older, dorsiflexion and plantarflexion of the ankle joints, and eversion and inversion in the range of motion also decrease with aging, and decline in range of motion has been identified as a risk factor of reduced balance ability²¹⁾. Thus, this study aimed to examine the influence of ankle range of motion on elderly people's balance ability in order to provide material for the prevention of falls and physical therapy interventions for enhancing balance ability.

SUBJECTS AND METHODS

This study randomly selected a total of 60 subjects, 30 old women and 30 old men aged 65 years or older, who gave consent to take part in this research. All subjects could walk independently. Exclusion criteria were a history of

surgery of the joints of the lower extremities, or taking drugs affecting balance ability. Before beginning the experiment, we explained the purposes of the research and experimental methods to the subjects.

TETRAX Portable Multiple System (TETRAX, Ramat Gan, and Sunlight Medical, Tel-Aviv, Israel) was used for the postural control test. The stability index indicates general stability by measuring the degree of postural sway at four sites on the foot, and it assesses an individual's ability to control and compensate postural changes. A higher stability index indicates greater instability^{22, 23)}, and frequent or greater changes of percentage of weight at the foot sites²⁴⁾. The synchronization index compares the waveforms of vibration measured at two foot sites of four body vibrations measured at each foot site, and a total of six combinations (AB, CD, AC, BD, AD, BC) are possible²⁵⁾.

Subjects stood barefoot on the foot plate holding a supporter and were tested in 8 postures for 32 sec. Movement for the test was limited and the test was started after confirming postural stabilization for 10 sec, not including data of the period of early stabilization. Then, the subjects were instructed to remove their hands from the supporter and to stand in an upright position. First, they were asked to stand on a stable base of support (hard plate) with their eyes open, and closed. Next they were asked to stand, with their eyes open and closed, on an unstable base of support (soft plate), which was created by putting a spring plate (31×12 cm) under the feet. When the subjects couldn't stand on the plate or couldn't keep an upright posture, a retest was conducted.

Measurement of ankle ROM was conducted by the method adopted by Norkin and White²⁶⁾ and the anatomical posture 0° was used as the initial posture for measuring the active range of motion. The range of motion was measured for dorsiflexion and plantarflexion of ankle joint and inversion and eversion. Dorsiflexion and plantarflexion were measured with a 180° stainless steel goniometer. The fixed arm was placed in parallel with the leg on the fibular side, the moving arm in parallel with foot on the 5th metatarsal side, and the axial point was placed on the lateral malleolus. Inversion and eversion were measured with an 8" (20 cm) goniometer. The fixed arm was placed in parallel with the leg on the tibia crest, the moving arm in parallel with the foot at the 2nd metatarsal, and the axial point at the mid-point between the medial malleolus and lateral malleolus. The test was performed by the same tester to minimize the difference between measurers.

Data measured in this study were statistically processed with SPSS version 12.0. General characteristics of the subjects were analyzed with descriptive statistics. Difference between the ankle angle and balance measurement were examined using the independent sample t-test. Correlations between ankle angle and balance ability were assessed using Pearson's correlation coefficient. A significance level α , of 0.05 was used in all analyses.

RESULTS

The comparison of RDF (right dorsiflexion), LDF

Table 1. Comparison of active range of motion by gender

	Male (n=30)	Female (n=30)
RDF	18.0 ± 4.0	17.6 ± 7.1
LDF	16.4 ± 4.8	16.7 ± 9.0
RPF	51.1 ± 10.5	45.7 ± 9.1*
LPF	49.8 ± 12.7	48.3 ± 9.9
RIN	12.7 ± 5.0	12.6 ± 5.6
LIN	13.8 ± 6.3	14.2 ± 7.1
REV	6.8 ± 3.1	5.8 ± 2.5
LEV	6.9 ± 4.2	5.1 ± 2.8*

(unit: degree) Mean ± SD. *p<0.05, RDF: Right dorsiflexion, LDF: Left dorsiflexion, RPF: Right plantarflexion, LPF: Light plantarflexion, RIN: Right inversion, LIN: Left inversion, REV: Right eversion, LEV: Left eversion

(left dorsiflexion), LPF (light plantarflexion), RIN (right inversion), LIN (left inversion), REV (right eversion) in active range of ankle motion showed no significant difference between male and female subjects. RPF (right plantarflexion) as 51.1 ± 10.5 for males, and 45.7 ± 9.1 for females, a significant difference (p<0.05). In addition, LEV (left eversion) was 6.9 ± 4.2 for males and 5.1 ± 2.8 for females, also a significant difference (p<0.05)(Table 1).

The comparison of stability index before and after visual deprivation on the stable base of support showed a tendency toward significant difference between male and female subjects, though the result itself was not significant (Table 2).

The comparison of the stability index before and after visual deprivation on the unstable plate showed no significant difference between male and female subjects. However, the stability index after visual deprivation was 49.7 ± 20.3 for men and 40.0 ± 12.7 for women, showing a statistically significant difference (p<0.05)(Table 3).

The comparison of the correlation of the stability index with active range of motion before and after visual deprivation on the stable base of support showed no significant difference in stability index before visual deprivation in active range of motion and stable base of support. However there were statistically significant differences in stability index after visual deprivation for male RPF (r=0.368) and LPF (r=0.395) (p<0.05)(Table 4).

DISCUSSION

As all motions of the ankle joints are related to maintaining balance in walking and control the interaction between the feet and the ground, they are vital for walking and balance¹⁸⁾. Thus, this study was designed to provide data necessary for physical therapy assessment and intervention programs. In this study we examined the balance ability of 60 out of 89 elderly people aged 65 years or older, through the, stability index and synchronization index measured before and after visual deprivation on an unstable base of support.

Vandercoort et al.²⁷⁾ reported that the range of motion

Table 2. Comparison of stability the index before and after visual deprivation on the stable base

	Male (n=30)	Female (n=30)
HOE (ST)	21.63 ± 9.13	17.99 ± 4.20
HCE (ST)	31.86 ± 14.55	26.54 ± 7.42

(n=60) HOE: Stable base, eyes open, HCE: Stable base, eyes Closed, ST: Stability

Table 3. Stability index before and after visual deprivation on the unstable base

	Male (n=30)	Female (n=30)
SOE (ST)	25.5 ± 8.6	24.1 ± 5.5
SCE (ST)	49.7 ± 20.4	40.0 ± 12.7*

(n=60) Mean ± SD. *p<0.05, SOE: Unstable base, eyes open, SCE: Unstable base, eyes closed, ST: Stability

Table 4. Correlations of stability index (ST) with active range of motion before and after visual deprivation on the stable base of support

		RDF	LDF	RPF	LPF	RIN	LIN	REV	LEV
HOE	Male (n=30)	-0.150	0.133	0.234	0.288	-0.165	0.076	0.030	-0.263
	Female (n=30)	0.233	0.175	0.050	0.088	0.024	0.051	-0.168	0.138
HCE	Male (n=30)	-0.092	0.106	0.368*	0.395*	-0.212	0.075	-0.105	-0.268
	Female (n=30)	0.243	0.105	0.024	0.260	-0.012	0.221	-0.284	0.091

(n=60) *p<0.05, HOE: Stable base, eyes open, HCE: Stable base, eyes closed, RDF: Right dorsiflexion, LDF: Left dorsiflexion, RPF: Right plantarflexion, LPF: Left plantarflexion, RIN: Right inversion, LIN: Left inversion, REV: Right eversion, LEV: Left eversion

tended to show more decrease in women than in men with increase of age, and Cunningham et al.²¹⁾ reported that decline in the range of motion was a risk factor related to decrease of balance ability. This study found there were significant differences from the respective male values of female RPF (right plantarflexion) and LEV (left eversion), and this finding is similar to those of Grimston, Vandercoort, and Cunningham.

Wilkerson et al.²⁸⁾ argued that as the height increase, balance ability decreases because of increase in body sway. The stability index before and after visual deprivation on a stable base of support was higher for males than for females. This means that the stability of their male subjects was lower, and the finding of this study was similar to Wilkerson's finding.

Lee and Kwon²⁹⁾ reported that there were significant changes in the balance index before and after visual deprivation on an unstable base and the balance index increased after visual deprivation. This means that COP sway was greater under visual deprivation, and that balance ability on an unstable base was reduced by visual deprivation. Thus, significant differences in the stability index due to after visual deprivation were similar to the findings of Lee and Kwon²⁹⁾.

Lee et al.³⁰⁾ reported that when static balance ability was assessed, forward and backward sway increased more than the left and right sway under the eyes open and closed conditions. Kwon and Choi³¹⁾ reported that subjects had difficulty with balance control under visual deprivation, exhibiting greater displacement from the reference point, and that the visual condition was an important factor in the maintenance of balance in an upright position regardless of the condition of the base of support. The active range of motion and male RPF (right plantarflexion) and LPF (left plantarflexion) showed significant changes after visual deprivation, and this finding is similar to the findings of Lee et al., and Kwon and Choi³¹⁾.

Research by Morgenroth et al.³²⁾ found that the balance

index differed between stable and unstable bases of support, increasing significantly as the base of support became unstable under all conditions. Our study also found significant changes in female LDF (left dorsiflexion) an unstable base of support, similar to the research of Morgenroth et al.³²⁾.

In the examination of correlations between active range of ankle motion and balance ability of the elderly people, we found significant difference in the results of TETRAX balance ability as the active range of motion increased. These findings explain that the active range of motion and visual deprivation affect balance ability. Therefore, we consider that a program for increasing the ankle range of motion of the elderly people, as a physical therapy intervention, would be beneficial for fall prevention and improve balance ability.

REFERENCES

- 1) Jeon MY: Effect of the fall prevention program (fpp) on gait, balance and muscle strength in elderly women at a nursing home. *J Korean Soc Biol Nurs Sci*, 2002, 4: 5-23.
- 2) Tinetti ME, Williams TF, Mayewski R: Fall risk index for elderly patients based on number of chronic disabilities. *Am J Med*, 1986, 80: 429-434. [Medline]
- 3) Wolfson LI, Whipple R, Amerman P, et al.: Stressing the postural response. A quantitative method for testing balance. *J Am Geriatr Soc*, 1986, 34: 845-850. [Medline]
- 4) Himann JE, Cunningham DA, Rechnitzer PA, et al.: Age-related changes in speed of walking. *Med Sci Sports Exerc*, 1988, 20: 161-166. [Medline]
- 5) Kerrigan DC, Todd MK, Della Croce U, et al.: Biomechanical gait alterations independent of speed in the healthy elderly: Evidence for specific limiting impairments. *Arch Phys Med Rehabil*, 1998, 79: 317-322. [Medline]
- 6) Gehlsen GM, Whaley MH: Falls in the elderly: Part ii, balance, strength, and flexibility. *Arch Phys Med Rehabil*, 1990, 71: 739-741. [Medline]
- 7) Mills EM: The effect of low-intensity aerobic exercise on muscle strength, flexibility, and balance among sedentary elderly persons. *Nurs Res*, 1994, 43: 207-211. [Medline]
- 8) Shumway-Cook A, Woollacott M, Kerns KA, et al.: The effects of two types of cognitive tasks on postural stability in older adults with and without a history of falls. *J Gerontol A Biol Sci Med Sci*, 1997, 52: M232-M240. [Medline]
- 9) Garbalosa JC, Cavanagh PR, Wu G, et al.: Foot function in diabetic patients after partial amputation. *Foot Ankle Int*, 1996, 17: 43-48. [Medline]

- 10) Lord SR, Clark RD, Webster IW: Postural stability and associated physiological factors in a population of aged persons. *J Gerontol*, 1991, 46: M69–M76. [[Medline](#)]
- 11) Song CH, Cha EJ, Lee TS, et al.: Balance plate to measure the center of pressure for evaluation of postural stability. *Chungbuk Med J*, 1995, 5: 61–69.
- 12) Chow DH, Cheng AC, Holmes AD, et al.: The effects of release height on center of pressure and trunk muscle response following sudden release of stoop lifting tasks. *Appl Ergon*, 2003, 34: 611–619. [[Medline](#)]
- 13) Chang MY, Lee YJ: The relationship between balance ability and visual perception ability in health elderly. *The Journal of Korean Academy of Occupational Therapy*, 1999, 7: 68–74.
- 14) Teasdale N, Bard C, LaRue J, et al.: On the cognitive penetrability of posture control. *Exp Aging Res*, 1993, 19: 1–13. [[Medline](#)]
- 15) Lord SR, Menz HB: Visual contributions to postural stability in older adults. *Gerontology*, 2000, 46: 306–310. [[Medline](#)]
- 16) Ronsky JL, Nigg BM, Fisher V: Correlation between physical activity and the gait characteristics and ankle joint flexibility of the elderly. *Clin Biomech (Bristol, Avon)*, 1995, 10: 41–49. [[Medline](#)]
- 17) Gong WT, Ma SY, Kim TH: The effect of ankle joint mobilization technique on equilibrium ability in the individuals with supinated foot. *Journal of the Korean Data And Information Science Society*, 2009, 20: 527–539.
- 18) Wolfson L, Whipple R, Judge J, et al.: Training balance and strength in the elderly to improve function. *J Am Geriatr Soc*, 1993, 41: 341–343. [[Medline](#)]
- 19) Woollacott MH, Shumway-Cook A, Nashner LM: Aging and posture control: Changes in sensory organization and muscular coordination. *Int J Aging Hum Dev*, 1986, 23: 97–114. [[Medline](#)]
- 20) Koon CH, Lee IH, Park KL: The effect of passive joint mobilization and massage on subacute lateral ankle ligament injuries. *The Journal Korean Society of Physical Therapy*, 2005, 17: 457–467.
- 21) Vandervoort AA, Chesworth BM, Cunningham DA, et al.: An outcome measure to quantify passive stiffness of the ankle. *Can J Public Health*, 1992, 83: S19–S23. [[Medline](#)]
- 22) Kohen-Raz R: Application of tetra-ataxiometric posturography in clinical and developmental diagnosis. *Percept Mot Skills*, 1991, 73: 635–656. [[Medline](#)]
- 23) Kohen-Raz R, Kohen-Raz A, Erel J, et al.: Postural control in pilots and candidates for flight training. *Aviat Space Environ Med*, 1994, 65: 323–326. [[Medline](#)]
- 24) Kim BR, Choi KH, Chun MH, et al.: Evaluation of balance control in patients with idiopathic parkinson's disease using tetra-ataxiometric posturography. *J Korean Acad Rehab Med*, 2009, 33: 538–546.
- 25) Kim CR, Chun MH, Lee GA: Assessments of balance control using tetra-ataxiometric posturography. *J Korean Acad Rehab Med*, 2009, 33: 429–435.
- 26) Norkin CC, White DJ: Measurement of joint motion: A guide to goniometry, third edition. *Physiotherapy Canada*, 2004, 56: 250.
- 27) Vandervoort AA, Chesworth BM, Cunningham DA, et al.: Age and sex effects on mobility of the human ankle. *J Gerontol*, 1992, 47: M17–M21. [[Medline](#)]
- 28) Wilkerson CG, King SM, Witman GB: Molecular analysis of the gamma heavy chain of chlamydomonas flagellar outer-arm dynein. *J Cell Sci*, 1994, 107: 497–506. [[Medline](#)]
- 29) Lee HS, Kwon HC: The influence of foot angle on standing balance upon the unstable platform. *Journal of the Korean Academy of University Trained Physical Therapists*, 1997, 4: 34–44.
- 30) Lee JW, Kwon OY, Yi CH, et al.: Effect of ankle strategy exercise on improvement of balance in elderly with impaired balance. *Korean J Health Promot Dis Prev*, 2008, 8: 158–167.
- 31) Kwon OY, Choi HS: Evaluation of the balance ability for 20 to 29 years old on the unstable platform. *Journal of the Korean Academy of University Trained Physical Therapists*, 1996, 3: 1–11.
- 32) Morgenroth DC, Shakir A, Orendurff MS, et al.: Low-back pain in trans-femoral amputees: Is there a correlation with static or dynamic leg-length discrepancy? *Am J Phys Med Rehabil*, 2009, 88: 108–113. [[Medline](#)]